

MCTP-02-22

April 2002

hep-th/0204231

New G_2 Metric, D6-branes and Lattice Universe

H. Lü

Michigan Center for Theoretical Physics

University of Michigan, Ann Arbor, Michigan 48109

ABSTRACT

We construct a new (singular) cohomogeneity-three metric of G_2 holonomy. The solution can be viewed as a triple intersection of smeared Taub-NUTs. The metric comprises three non-compact radial-type coordinates, with the principal orbits being a T^3 bundle over S^1 . We consider an M-theory vacuum $(\text{Minkowski})_4 \times \mathcal{M}_7$ where $\mathcal{M}_7$ is the G_2 manifold. Upon reduction on a circle in the T^3 , we obtain the intersection of a D6-brane, a Taub-NUT and a 6-brane with R-R 2-form flux. Reducing the solution instead on the base space S^1 , we obtain three intersecting 6-branes all carrying R-R 2-form flux. These two configurations can be viewed as a classical flop in the type IIA string theory. After reducing on the full principal orbits and the spatial world-volume, we obtain a four-dimensional metric describing a lattice universe, in which the three non-compact coordinates of the G_2 manifold are identified with the spatial coordinates of our universe.

1 Introduction

Seven-dimensional manifolds of G_2 holonomy have long been known to exist. The construction of explicit non-compact G_2 metrics began ten years ago, when asymptotically conical metrics of cohomogeneity one were found [1, 2]. The physical interest of G_2 manifolds has increasingly significantly with the discovery of M-theory, because they are the most natural compactifying spaces from the eleven-dimensional point of view. It is expected that M-theory compactified on a G_2 manifold gives rise to an $\mathcal{N} = 1$ super Yang-Mills theory in $D = 4$ [3]. The G_2 manifold with principal orbits $S^3 \times S^3$ provides a geometrical demonstration of the classical flop of the type IIA superstring theory [4]. In [5], M-theory dynamics on a G_2 manifold were discussed.

Recently, a large class of new metrics of G_2 holonomy have been obtained [6]-[17], following the construction of the first examples of asymptotically locally conical spin(7) manifolds [18]. These examples have non-abelian isometry groups. G_2 metrics with nilpotent isometry groups were also constructed in [19], which can be obtained by taking the Heisenberg or Euclidean limits of the non-abelian examples. Whilst it is of great interest to construct regular G_2 metrics, physically, it is essential to have an appropriate singularity structure to give rise to chiral fermions in $D = 4$ [20, 21].

In section 2, we construct a new non-compact cohomogeneity three metric with G_2 holonomy. The metric has three non-compact radial-type coordinates, with the principal orbits being a T^3 bundle over S^1 . The isometry group of the metric is a four-dimensional nilpotent Lie group. The metric has either power-law singularities or delta-function singularities. The solution can be viewed as the intersection of three smeared Taub-NUTs. When one of the Taub-NUT charges is set to zero, the metric describes a product of S^1 with a six-dimensional non-compact Calabi-Yau manifold.

In section 3, we consider an M-theory vacuum $(\text{Minkowski})_4 \times \mathcal{M}_7$, where $\mathcal{M}_7$ is the G_2 manifold. We show that by dimensionally reducing the solution on one of the circles in the T^3 , we obtain a type IIA configuration with one D6-brane, one Taub-NUT and one 6-brane with an R-R 2-form flux. On the other hand, if we reduce the solution on the base space S^1 , we obtain an intersection of three 6-branes all carrying R-R 2-form flux. These two configurations can be viewed as the classical flop in type IIA string theory on a non-compact six-dimensional Kähler manifold with a nilpotent isometry group. The origin of the flop is that the T^3 bundle over S^1 principal orbits can also be viewed as S^1 bundle over T^3 .

In section 4, we perform a Kaluza-Klein reduction on the full principal orbits and the

spatial world-volume. We obtain three perpendicularly intersecting membranes in $D = 4$, describing a lattice universe. In this picture, the three non-compact coordinates of the G_2 manifold are identified with the spatial coordinates of our universe. We conclude the letter in section 5.

2 New G_2 metric

The metric ansatz is given by

$$\begin{aligned} ds_7^2 = & H_1 dx_1^2 + H_2 dx_2^2 + H_3 dx_3^2 + H_1 H_2 H_3 dz_4^2 + H_3 H_1^{-1} (dz_1 + H_1' z_2 dz_4)^2 \\ & + H_1 H_2^{-1} (dz_2 + H_2' z_3 dz_4)^2 + H_2 H_3^{-1} (dz_3 + H_3' z_1 dz_4)^2, \end{aligned} \quad (1)$$

where H_1 , H_2 and H_3 are functions of x_1 , x_2 and x_3 respectively. The prime on H_i' denotes a derivative with respect to the argument of H_i ;

$$H_1' = \partial_{x_1} H_1, \quad H_2' = \partial_{x_2} H_2, \quad H_3' = \partial_{x_3} H_3. \quad (2)$$

The natural vielbein basis is

$$\begin{aligned} e^0 &= \sqrt{H_1 H_2 H_3} dz_4, & e^1 &= \sqrt{H_1} dx_1, & e^2 &= \sqrt{H_2} dx_2, & e^3 &= \sqrt{H_3} dx_3, \\ e^4 &= \sqrt{H_3 H_1^{-1}} (dz_1 + H_1' z_2 dz_4), & e^5 &= \sqrt{H_1 H_2^{-1}} (dz_2 + H_2' z_3 dz_4), \\ e^6 &= \sqrt{H_2 H_3^{-1}} (dz_3 + H_3' z_1 dz_4). \end{aligned} \quad (3)$$

The associative 3-form in this basis is given by

$$\Phi = e^{016} + e^{024} + e^{035} + e^{125} - e^{134} + e^{236} - e^{456}, \quad (4)$$

where $e^{ijk} = e^i \wedge e^j \wedge e^k$. The metric (1) has G_2 holonomy if and only if Φ is closed and co-closed. We find that the closure and co-closure of Φ implies that

$$H_i'' = 0, \quad i = 1, 2, 3, \quad (5)$$

implying that

$$H_1 = 1 + m_1 x_1, \quad H_2 = 1 + m_2 x_2, \quad H_3 = 1 + m_3 x_3. \quad (6)$$

Here the constant 1 is included so that H_i does not vanish when $m_i = 0$. Clearly the metric has a power-law singularity whenever any of the H_i vanishes. The metric can also be recast in a “co-moving” frame,

$$\begin{aligned} ds_7^2 = & dr_1^2 + dr_2^2 + dr_3^2 + \frac{9}{4} (m_1 m_2 m_3 r_1 r_2 r_3)^{2/3} dz_4^2 + \left(\frac{m_3 r_3}{m_1 r_1} \right)^{2/3} (dz_1 + m_1 z_2 dz_4)^2 \\ & + \left(\frac{m_1 r_1}{m_2 r_2} \right)^{2/3} (dz_2 + m_2 z_3 dz_4)^2 + \left(\frac{m_2 r_2}{m_3 r_3} \right)^{2/3} (dz_3 + m_3 z_1 dz_4)^2. \end{aligned} \quad (7)$$

The fibration in the z_1 , z_2 and z_3 coordinates implies that the constants m_i are quantised, namely [22]

$$m_1 = n_1 \frac{L_1}{L_2 L_4}, \quad m_2 = n_2 \frac{L_2}{L_3 L_4}, \quad m_3 = n_3 \frac{L_3}{L_1 L_4}, \quad (8)$$

where n_i are integers and L_i are the periods of the z_i . For simplicity, we can set $L_i = \ell_p$ where ℓ_p is the Plank length, and then $m_i = n_i/\ell_p$.

The metric has a power-law singularity when any of the H_i vanishes. This can be avoided by instead taking

$$H_i = 1 + \sum_{\alpha} m_i^{\alpha} |x_i - x_i^{\alpha}|. \quad (9)$$

such that H_i is positive definite. However, in doing so, we have introduced delta function singularities.

When all three of the H'_i are non-vanishing, the metric describes three intersecting Taub-NUTs with three independent non-vanishing smeared charges. The metric has three non-compact coordinates x_1 , x_2 and x_3 . The principal orbits are T^3 bundle over S^1 ; they are parameterised by the coordinates (z_1, z_2, z_3) and z_4 respectively. The metric is of co-homogeneity three since it depends explicitly on the three non-compact coordinates x_i . Despite the dependence on the (z_1, z_2, z_3) coordinates, they, together with z_4 , parameterise a four-dimensional nilpotent Lie group $\mathcal{G}$, which is the isometry group of the metric, and thus the four-dimensional principal orbits are homogeneous.

When two of the H'_i vanish, the metric describes a direct product of Euclidean 3-space and a smeared Taub-NUT. If instead only one of H'_i vanishes, in which case the metric was obtained in [23], it describes a product of an S^1 with a Calabi-Yau 6-manifold. To see this in detail, let us set $H_3 = 1$. The metric of the Calabi-Yau manifold is then given by

$$\begin{aligned} ds_6^2 = & H_1 dx_1^2 + H_2 dx_2^2 + H_1 H_2 dz_4^2 + H_2 dz_3^2 \\ & + H_1^{-1} (dz_1 + H'_1 z_2 dz_4)^2 + H_1 H_2^{-1} (dz_2 + H'_2 z_3 dz_4)^2. \end{aligned} \quad (10)$$

and the Kähler form is given by

$$J = e^0 \wedge e^5 - e^1 \wedge e^4 + e^2 \wedge e^6, \quad (11)$$

where the vielbein is given by (3) with $H_3 = 1$.

3 Intersecting D6-branes

Having obtained the new G_2 metric, one may consider an M-theory vacuum solution given by the direct product of Minkowski 4-spacetime and the G_2 manifold, namely

$$ds_{11}^2 = -dt^2 + dw_1^2 + dw_2^2 + dw_3^2 + ds_7^2. \quad (12)$$

The solution can be viewed as a triple intersection of smeared Taub-NUTs, with the metric represented by the diagram

	t	w_1	w_2	w_3	x_1	x_2	x_3	z_1	z_2	z_3	z_4
$H_1(x_1)$	$\times$	$\times$	$\times$	$\times$	$-$	$\times$	$\times$	$*$	$-$	$\times$	$-$
$H_2(x_2)$	$\times$	$\times$	$\times$	$\times$	$\times$	$-$	$\times$	$\times$	$*$	$-$	$-$
$H_3(x_3)$	$\times$	$\times$	$\times$	$\times$	$\times$	$\times$	$-$	$-$	$\times$	$*$	$-$

Diagram 1. Triple intersections of Taub-NUTs. Here $\times$, $-$ and $*$ denote the world-volume, transverse space, and fibre coordinates respectively.

There is a $U(1)$ isometry for each of the z_i coordinates, and so we can reduce the metric on any z_i to obtain a solution in type IIA theory. The z_i for $i = 1, 2, 3$ are equivalent, and hence the reduction can be discussed using z_1 as a representative. The resulting type IIA solution is given by

$$\begin{aligned}
e^{\frac{1}{6}\phi} ds_{10}^2 &= -dt^2 + dw_1^2 + dw_2^2 + dw_3^2 + H_1 dx_1^2 + H_2 dx_2^2 + H_3 dx_3^2 + H_1 H_2 H_3 dz_4^2 \\
&\quad + H_1 H_2^{-1} (dz_2 + H_2' z_3 dz_4)^2 + H_2 H_3^{-1} dz_3^2 + H_3 H_1 (H_1' z_2)^2 dz_4^2 \\
&\quad - W^{-1} (H_3 H_1^{-1} H_1' z_2 dz_4 - H_2 H_3^{-1} H_3' z_4 dz_3)^2, \\
e^\phi &= W^{\frac{3}{4}}, \quad W = H_3 H_1^{-1} + H_2 H_3^{-1} (H_3' z_4)^2, \\
A_{(1)} &= W^{-1} (H_3 H_1^{-1} H_1' z_2 dz_4 - H_2 H_3^{-1} H_3' z_4 dz_3). \quad (13)
\end{aligned}$$

Note that before performing the Kaluza-Klein reduction, we have made a coordinate transformation $z_3 \rightarrow z_3 - H_3' z_1 z_4$ in the metric (1). Clearly, the solution describes an intersection of three objects. The one parameterised by H_1 is a smeared D6-brane, and the one parameterised by H_2 is a Taub-NUT. The one associated with H_3 is a 6-brane carrying an R-R 2-form flux, but it differs from a standard D6-brane.

We can instead reduce the solution on the z_4 coordinate, giving the type IIA solution

$$e^{\frac{1}{6}\phi} ds_{10}^2 = -dt^2 + dw_1^2 + dw_2^2 + dw_3^2 + H_1 dx_1^2 + H_2 dx_2^2 + H_3 dx_3^2$$

$$\begin{aligned}
& +H_3 H_1^{-1} dz_1^2 + H_1 H_2^{-1} dz_2^2 + H_2 H_3^{-1} dz_3^2 \\
& -W^{-1} (H_3 H_1^{-1} H'_1 z_2 dz_1 + H_1 H_2^{-1} H'_2 z_3 dz_2 + H_2 H_3^{-1} H'_3 z_1 dz_3)^2, \\
e^\phi = W^{\frac{3}{4}}, \quad W = H_1 H_2 H_3 + H_3 H_1^{-1} (H'_1 z_2)^2 + H_1 H_2^{-1} (H'_2 z_3)^2 + H_2 H_3^{-1} (H'_3 z_1)^2, \\
A_{(1)} = W^{-1} (H_3 H_1^{-1} H'_1 z_2 dz_1 + H_1 H_2^{-1} H'_2 z_3 dz_2 + H_2 H_3^{-1} H'_3 z_1 dz_3). \tag{14}
\end{aligned}$$

The solution describes three intersecting 6-branes all carrying R-R 2-form flux. These 6-branes are different from the usual D6-brane coming from the reduction of the fibre coordinate of a Taub-NUT in $D = 11$.

The two configurations arising from the reduction on z_1 or z_4 can be viewed as a classical flop in the type IIA string theory on the non-compact Kähler manifold. The flop in $D = 10$ can be geometrically explained by the fact that the T^3 bundle over S^1 principal orbits of the four-dimensional nilpotent Lie group $\mathcal{G}$ can also be described as an S^1 bundle over T^3 . However, the two descriptions are somewhat different. In the latter case, the fibre is the circle group in $\mathcal{G}$ generated by $\frac{\partial}{\partial z_4}$. In the former case, the fibre is not the orbit of a three-dimensional subgroup of $\mathcal{G}$ because $\frac{\partial}{\partial z_i}$ for $i = 1, 2, 3$ are not themselves Killing vectors; we must add a multiple of $\frac{\partial}{\partial z_4}$. In fact the flop involves interchanging the fibre and base spaces of the $U(1)$ fibration. This is analogous to the flop discussed in [4].

4 Lattice universe

The new G_2 metric (1) that we have obtained is in fact inspired by the four-dimensional intersecting membrane solution that describes the lattice universe [23].¹ There has been experimental evidence suggesting that the network of galaxy superclusters and voids seems to form a three-dimensional lattice with a spacing of about $120h^{-1} \text{ Mpc}$ (where h^{-1} is the Hubble constant in units of $100 \text{ km s}^{-1} \text{ Mpc}^{-1}$) [25, 26, 27]. In [23], an M-theory solution was constructed to describe such a lattice structure, which can be realised by considering non-standard brane intersections of two M5-branes and one Taub-NUT, or two Taub-NUTs and one M5-brane. In the latter case, turning off the M5-brane charge causes the solution to reduce to the product of 5-dimensional Minkowski spacetime and the non-compact Calabi-Yau manifold given in (10).

In section 2, we obtained the new G_2 metric (1) by adding an extra fibration on the seventh coordinate. This procedure follows the general prescription of obtaining G_2 manifolds from six-dimensional Kähler manifolds, described in detail in [28].

¹In [24], a triply quasi-periodic Gibbons-Hawking metric was obtained.

If we reduce the M-theory solution on the world-volume spatial coordinates w_i and also on the T^3 bundle over S^1 principal orbits, we obtain three perpendicularly intersecting membranes in $D = 4$, with the metric

$$ds_4^2 = (H_1 H_2 H_3)^{\frac{1}{2}} (-dt^2 + H_1 dx_1^2 + H_2 dx_2^2 + H_3 dx_3^2). \quad (15)$$

This metric was first obtained in [23], although it was supported by very different field strength. The functions of H_i in this case are given by (9) describing periodic arrays of intersecting membranes.

In this static cosmological model the spatial world-volume and the T^3 bundle over S^1 principal orbits are viewed as an internal space, whilst the three non-compact coordinates x_1 , x_2 and x_3 of the G_2 manifold are identified with the spatial coordinates of our universe. This is rather natural since the principal orbits are clearly compact, and the spatial world-volume can be wrapped on a compact space such as T^3 . Although it is not likely that the metric (15) describes our actual universe, since it preserves $\mathcal{N} = 1$ supersymmetry; it is nevertheless rather suggestive that the lattice structure should emerge from a metric with G_2 holonomy.

5 Conclusions

In this letter, we constructed a new cohomogeneity-three metric with G_2 holonomy. It has three radial-type coordinates, with the principle orbits being a T^3 bundle over S^1 . The solution can be viewed as three intersecting Taub-NUTs. We performed Kaluza-Klein reduction on the S^1 and instead on a circle in the T^3 . The two resulting type IIA configurations can be viewed as a classical flop in type IIA string theory on a non-compact Kähler six-manifold. Although the type IIA solutions do not describe the triply intersecting D6-branes advocated in [21, 29] for the realisation of chiral fermions, it is nevertheless of interest to investigate further if chirality could arise from the singularities of our G_2 metric. The metric provides a concrete example for studying such an issue, since it can be viewed as the lifting of a D6-brane configurations in $D = 10$.

If we perform a Kaluza-Klein reduction on the entire four-dimensional full principal orbits and the spatial world-volume, we obtain triply intersecting membranes in $D = 4$, describing a lattice universe. The construction takes full advantage of the non-compact nature of the manifold, in that the three non-compact coordinates are precisely identified with the spatial coordinates of our universe. It is of great interest to investigate further the significance of such a configuration arising from a G_2 manifold.

Acknowledgement

We are grateful to Mirjam Cvetič for extensive discussions on triply intersecting D6-branes and chiral fermions, to Gary Gibbons for pointing out the isometry group of the metric, and to Chris Pope and Justin Vázquez-Poritz for discussions.

References

- [1] R.L. Bryant and S. Salamon, *On the construction of some complete metrics with exceptional holonomy*, Duke Math. J. **58**, 829 (1989).
- [2] G.W. Gibbons, D.N. Page and C.N. Pope, *Einstein metrics on S^3 , $\mathbb{R}^3$ and $\mathbb{R}^4$ bundles*, Commun. Math. Phys. **127**, 529 (1990).
- [3] B.S. Acharya, *On realising $\mathcal{N} = 1$ super Yang-Mills in M theory*, hep-th/0011089.
- [4] M. Atiyah, J. Maldacena and C. Vafa, *An M-theory flop as a large N duality*, J. Math. Phys. **42**, 3209 (2001), hep-th/0011256.
- [5] M. Atiyah and E. Witten, *M-theory dynamics on a manifold of G_2 holonomy*, hep-th/0107177.
- [6] M. Cvetič, G.W. Gibbons, H. Lü and C.N. Pope, *Supersymmetry M2-branes and G_2 manifolds*, Nucl. Phys. **B620**, 3 (2002), hep-th/0106026.
- [7] A. Brandhuber, J. Gomis, S.S. Gubser and S. Gukov, *Gauge theory at large N and new $G(2)$ holonomy metrics*, Nucl. Phys. **B611**, 179 (2001), hep-th/0106034.
- [8] N. Hitchin, *Stable forms and special metrics*, math.DG/0107101.
- [9] M. Cvetič, G.W. Gibbons, H. Lü and C.N. Pope, *Cohomogeneity one manifolds of $Spin(7)$ and G_2 holonomy*, hep-th/0108245.
- [10] M. Cvetič, G.W. Gibbons, H. Lü and C.N. Pope, *Orientifolds and slumps in G_2 and $Spin(7)$ metrics*, hep-th/0111096.
- [11] M. Cvetič, G.W. Gibbons, H. Lü and C.N. Pope, *M-theory conifolds*, Phys. Rev. Lett. **88**, 121602 (2002), hep-th/0112098.
- [12] A. Brandhuber, *G_2 holonomy spaces from invariant three-forms*, hep-th/0112113.

- [13] M. Cvetič, G.W. Gibbons, H. Lü and C.N. Pope, *A G_2 unification of the deformed and resolved conifolds*, hep-th/0112138, to appear in Phys. Lett. **B**.
- [14] R. Hernandez and K. Sfetsos, *An eight-dimensional approach to G_2 manifolds*, hep-th/0202135.
- [15] S. Gukov, S-T. Yau and E. Zaslow, *Duality and fibrations on G_2 manifolds*, hep-th/0203217.
- [16] K. Behrndt, *Singular 7-manifolds with G_2 holonomy and intersecting 6-branes*, hep-th/0204061.
- [17] Z.W. Chong, M. Cvetič, G.W. Gibbons, H. Lü, C.N. Pope and P. Wagner, *General metrics of G_2 holonomy and contraction limits*, hep-th/0204064.
- [18] M. Cvetič, G. W. Gibbons, H. Lü and C. N. Pope, *New complete non-compact $Spin(7)$ manifolds*, hep-th/0103155; *New cohomogeneity one metrics with $Spin(7)$ holonomy*, math.DG/0105119.
- [19] G.W. Gibbons, H. Lü, C.N. Pope and K.S. Stelle, *Supersymmetric domain walls from metrics of special holonomy*, Nucl. Phys. **B623**, 3 (2002), hep-th/0108191.
- [20] B. Acharya and E. Witten, *Chiral fermions from manifolds of G_2 holonomy*, hep-th/0109152.
- [21] M. Cvetič, G. Shiu and A.M. Uranga, *Chiral type II orientifold constructions as M theory on G_2 holonomy spaces*, hep-th/0111179.
- [22] I.V. Lavrinenko, H. Lü and C.N. Pope, *Fibre bundles and generalised dimensional reductions*, Class. Quant. Grav. **15**, 2239 (1998), hep-th/9710243.
- [23] M.J. Duff, P. Hoxha, H. Lü, R.R. Martinez-Acosta and C.N. Pope, *A lattice universe from M -theory*, Phys. Lett. **B451**, 38 (1999), astro-ph/9712301.
- [24] S. Nergiz and C. Saclioglu, *A Quasiperiodic Gibbons-Hawking metric and space-time foam*, Phys. Rev. **D53**, 2240 (1996), hep-th/9505141.
- [25] J. Einasto, M. Einasto, S. Gottlober, V. Muller, V. Saar, A.A. Starobinsky, E. Tago, D. Tucker, H. Andernach and P. Frisch, *A 120 Mpc periodicity in the three-dimensional distributions of galaxy superclusters*, Nature, **385** (1997) 139.

- [26] R. Kirshner, *The universe as a lattice*, Nature, **385** (1997).
- [27] J. Einasto, *Has the universe a honeycomb structure*, astro-ph/9711320.
- [28] M. Cvetič, G.W. Gibbons, H. Lü and C.N. Pope, *Almost special holonomy in type IIA and M theory*, hep-th/0203060.
- [29] E. Witten, *Deconstruction, G_2 holonomy, and doublet-triplet splitting*, hep-ph/0201018.